

# Berries

iNvigate® | B Sure® | UpLift®

## PRODUCT APPLICATION AND TIMING FOR BERRIES (SUCH AS RED AND BLACK CANEBERRIES, BLUEBERRIES AND STRAWBERRIES)

### B Sure®

- Supports plant productivity in a variety of environmental conditions
- Contains carbon, nitrogen, true protein, amino acids and essential metabolic micronutrients

### iNvigate®

- Highly productive microbial community when applied to the soil, assisting in nutrient uptake and improving fertilizer efficiency
- Promotes availability and absorption of nitrogen, phosphorus and potassium. Along with promoting a healthy soil microbial community, also promotes better fertilizer utilization and increased yield

### UpLift®

- Fertility solution derived from organic, biologically-extracted chitin and additional critical nutrients
- Promotes robust soil microbial environment and interacts with the plant to increase crop productivity

### B Sure®

Type Of Application	pt/A	Timing
Foliar	1 - 4	Every 2 to 4 weeks <sup>1</sup>
Soil	1 - 4	Every 2 to 4 weeks <sup>2</sup>

<sup>1</sup>With foliar nutrition, fungicides, herbicides and insecticides from pre-bloom through harvest.

<sup>2</sup>From transplant to 60 days after transplant.

### iNvigate®

Type Of Application	pt/A	Timing
Soil	2 - 4	Every 3 to 4 weeks <sup>3</sup>

<sup>3</sup>With fertilizers, micronutrients, fungicides, herbicides or insecticides from planting or transplant through harvest.


### UpLift®

Type Of Application	lb/A	Timing
Soil	0.5 - 2	Every 3 to 4 weeks <sup>3</sup>


<sup>3</sup>With fertilizers, micronutrients, fungicides, herbicides or insecticides from planting or transplant through harvest.

\*AMVAC products not sold in all U.S. States.

Foliar Application Timing



Soil Application Timing



Visit our website at [www.AMVAC.com](http://www.AMVAC.com)