

POSTEMERGENCE APPLICATION GUIDELINES IN CORN

Active Ingredient	<ul style="list-style-type: none"> Topramezone: 2.8 pounds of topramezone free acid per gallon.
Labeled Uses	<ul style="list-style-type: none"> All types* of field corn, popcorn, sweet corn and between crop applications. *Including inbreds. Refer to seed company recommendations before application of Impact® Herbicide on inbred lines.
Recommended Rate	<ul style="list-style-type: none"> 0.75–1.0 fl oz/A (0.016–0.022 lb topramezone), see the reverse page for use rates on specific weeds.
Tank Additives for Best Performance	<ul style="list-style-type: none"> AMS at 2–2.5 lb/A or UAN 28–34% at 3 pt/A. <ul style="list-style-type: none"> Provides ammonium ion transport of weak acid herbicides to site of action. Liquid AMS (3.4 lb AMS/gal) at 5–6 pt/A may be substituted. MSO at 1–1.5% v/v (1–1.5 gal/100 gal of water), containing minimum 10% emulsifier. <ul style="list-style-type: none"> Provides more effective leaf cuticular wax penetration than other adjuvants. Use higher rate during periods of hot, dry weather and to larger weeds. With glyphosate tank mixtures, use 0.5% v/v (2 qt/100 gal of water) and AMS.
Tank Mix Partners	<ul style="list-style-type: none"> Atrazine is recommended at 0.25–1.0 lb/A. For a second effective mode of action, 1 lb/A minimum is recommended. Other partners include glyphosate, glufosinate, Status®, DiFlexx® and residual corn herbicides.
Application Window	<ul style="list-style-type: none"> Optimum timing is 2- to 4-inch height, weed species dependent. Impact® Herbicide may be applied up to 45 days before harvest.
Application Guidelines	<ul style="list-style-type: none"> Recommended water volume: Apply in 15 gal/A by ground and 4 gal/A by air to obtain thorough coverage. For ground applications, flat-fan nozzles delivering medium droplets (250–400 micron) are recommended. Drop nozzles and/or higher spray volume are recommended if the crop canopy prevents adequate spray coverage of weeds. Application should be made a minimum of one hour before rainfall or irrigation.
Insecticide Guidelines	<ul style="list-style-type: none"> Impact Herbicide may be safely applied following a soil-applied organophosphate insecticide, such as COUNTER 20G, or tank-mixed with a foliar-applied insecticide.

Impact Tank-Mixing Order Recommendation

1. Fill spray tank $\frac{1}{2}$ to $\frac{3}{4}$ full with clean water.
2. Add dry or liquid nitrogen fertilizers (AMS or UAN).
3. Add **Impact Herbicide**.
4. Add WP (wettable powder), DG (dispersible granule), DF (dry flowable) or F (liquid flowable) formulations.
5. Add EC (emulsifiable concentrate) formulations.
6. Add spray adjuvants (MSO).
7. Fill the remainder of the tank with water.
(Refer to label for full tank mix directions for use)

Impact Aerial Application Guidelines

- AMS at 1 lb/A or UAN at 1 qt/A
- plus**
- Impact** at 0.75 –1.0 fl oz/A
- plus**
- Atrazine 4L at 0.5–1.0 pt/A (if corn height allows)
- plus**
- Methylated seed oil (MSO) at 0.5 pt/A
- Application volume of 4 gal/A recommended for thorough coverage
 - Approved drift-reducing additive may be added in the tank-mix after MSO

Impact – Postemergence Application Guidelines in Corn

Broadleaf Weeds Controlled by Impact at 0.75–1.0 fl oz/A and Maximum Recommended Weed Height in Inches at Application¹

Annual Broadleaf Weeds ²	Maximum Size (in) ³ 0.75–1.0 fl oz/A	Annual Broadleaf Weeds ²	Maximum Size (in) ³ 0.75–1.0 fl oz/A
Amaranth, Palmer	6 ⁴	Nightshade, Hairy	6
Amaranth, Powell	6	Pigweed, Prostrate	6
Burcucumber	6	Pigweed, Redroot	6
Canola, Volunteer	6	Pigweed, Smooth	6
Carpetweed	6	Pigweed, Tumble	4
Chickweed, Common	4	Prickly Lettuce	4
Cocklebur, Common	8	Pusley, Florida	3
Dandelion	6 ⁵	Ragweed, Common	6
Galinsoga, Hairy	6	Ragweed, Giant	8 ⁴
Henbit	4	Shepherd's-purse	4
Jimsonweed	6	Sida, Prickly	3
Kochia	6 ⁴	Smartweed, Ladysthumb	3
Lambsquarters, Common	6	Smartweed, Pennsylvania	3
Mallow, Common	3	Sunflower, Volunteer	8 ⁴
Mallow, Venice	3 ⁵	Sunflower, Wild (Common)	8 ⁴
Marestail (Horseweed)	6 ⁴	Thistle, Canada	6 ⁵
Morningglories	6 ⁵	Thistle, Russian	4
Mustards	6	Velvetleaf	8 ⁴
Nightshade, Black	6	Waterhemp, Common	6 ⁴
Nightshade, Eastern Black	6	Waterhemp, Tall	6 ⁴

¹Refer to EPA label for information on best management and stewardship practices regarding herbicide resistance.

²Refer to Section X of the EPA label for list of scientific names.

³For best performance, apply before weeds exceed the maximum size or vine length listed.

⁴A rate of 1.0 fl oz/A is recommended if maximum weed size is reached.

⁵Indicates growth suppression.

Grass Weeds Controlled by Impact at 0.75 or 1.0 fl oz/A and Maximum Recommended Weed Height in Inches at Application¹

Annual Grass Weeds ^{2,3}	Impact at 0.75 fl oz/A	Impact at 1.0 fl oz/A	Annual Grass Weeds ^{2,3}	Impact at 0.75 fl oz/A	Impact at 1.0 fl oz/A
Barnyardgrass	4	5	Goosegrass	3	4
Crabgrass, Large	3	4	Johnsongrass, Seedling	4 ⁴	4
Crabgrass, Smooth	3	4	Millet, Wild-Proso	3	4
Cupgrass, Woolly	3 ⁴	3	Panicum, Fall	3 ⁴	3
Foxtail, Giant	4	5	Panicum, Texas	3 ⁴	3 ⁴
Foxtail, Green	3	4	Shattercane	4 ⁴	4 ⁴
Foxtail, Yellow	3 ⁴	3	Signalgrass, Broadleaf	3 ⁴	3

¹Refer to EPA label for information on best management and stewardship practices regarding herbicide resistance.

²Refer to Section X of the EPA label for list of scientific names.

³For best performance, apply before grasses exceed the maximum size listed in this table.

⁴Partial control of grass weeds with Impact at rates shown.

See our entire line of products at AMVAC.com



©2020 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All rights reserved. AMVAC, Counter, Impact and respective logos are trademarks owned by AMVAC Chemical Corporation. Status is a trademark of BASF Corporation. DiFlexx is a trademark of Bayer Intellectual Property.

Atrazine and Counter are restricted use pesticides. Important: Always read and follow label instructions. See label for listed weeds and pests. Some products may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status. All products are EPA registered. A290-064504 9/20