

POSTEMERGENCE BROADLEAF WEED CONTROL IN FIELD CORN

Product Advantages

- The preferred tank-mix partner for control of hard-to-manage and glyphosate-resistant broadleaf weeds.
- Use on conventional and all genetically traited field corn.
- Wide application window: Preplant to V6 (6 collar), 20-inch corn.
- HRAC Group 2 and Group 4 modes of action for resistance management and maximum performance.
- Low concern for auxin herbicide volatility.
- Excellent corn tolerance.
- Does not contain atrazine or 2,4-D.

Annual Weeds Controlled:

Cocklebur, Henbit, Horseweed, Jimsonweed, Mustard spp., Nightshade spp., Prickly Sida, Purslane, Common and Giant Ragweed, Sicklepod, Annual Spurge spp., Smartweed spp., Sunflower, Velvetleaf, Annual/Biennial Wormwood

Biennial/Perennial Weeds Controlled:

Volunteer Alfalfa, Jerusalem Artichoke, Burdock, Clover, Curly Dock, Dandelion, Red Sorrel

Weeds Partially Controlled:

Canada Thistle, Kochia¹, Lambsquarters¹, Annual Morningglory spp., Pigweed spp¹, Russian Thistle¹, Wild Buckwheat

¹Will not control ALS-resistant weeds.

Application Timing

- Preplant, preemergence and postemergence up to V6 (6 collars) and 20-inch tall corn.
- Apply with recommended adjuvants when emerged weeds are present at application.
- Apply with products containing acetochlor, atrazine, dicamba, mesotrione, metolachlor or topramezone for complete control of grass and broadleaf weeds and to increase residual weed control.
- Apply postemergence with glyphosate to glyphosate-resistant field corn varieties.

Crop Rotation Interval

Barley, Oat, Rye, Wheat	4 Months
Alfalfa ¹ , Cover Crops ¹ , Dry Beans ¹ , Lima Beans, Forage Grasses ¹ , Popcorn, Rice, Soybean ¹	10.5 Months
Grain Sorghum	12 Months
Snap Bean ¹ , Sweet Corn ¹ , Cotton, Pea ¹ , Peanut, Potato, Sunflower, Tobacco	18 Months
Canola, Sugarbeet and All Other Crops	26 Months and Bioassay

¹Refer to product label for restrictions and different rotation intervals.

Mixing Instructions

- 2–5 oz/A (refer to product label for weeds controlled at different rates).
- Use a spray volume of 10–40 gal/A.
- For postemergence, applications add the following adjuvants: NIS at 0.25% v/v, COC at 1% v/v, or MSO at 1% v/v plus AMS at 2–4 lbs/A or UAN at 2.5% v/v.
- Do not use undiluted UAN as total carrier.

Other Information

- Rainfast in 2 hours.
- Aerial application not allowed.
- Do not apply to sweet corn or popcorn.
- Preharvest interval is 85 days for field corn grown for grain and 45 days for field corn grown for forage or ensilage.
- Maximum total application rate is 6 oz/A in a single crop year.
- Do not apply **Hornet**[®] Herbicide if corn was previously treated by the organophosphate insecticides, COUNTER[®] Insecticide or Thimet[®] Insecticide.

See our entire line of products at AMVAC.com

Hornet Herbicide
HRAC Group 2, 4