

ASSURE® II HERBICIDE CONTROLS TOUGH GRASSES AND VOLUNTEER CORN IN SOYBEAN

Assure® II Herbicide Delivers:

- Controls volunteer conventional (non-trait), Roundup Ready® or LibertyLink® trait containing corn, but not quizalofop resistant corn (e.g., Enlist® corn).
- Controls other grass species such as fall panicum, field sandbur, giant foxtail, johnsongrass, shattercane, broadleaf singalgrass, barnyardgrass, sprangletop and quackgrass.
- **Assure II®** can be tank-mixed with several broadleaf herbicides and insecticides.
- Approved tank-mix partner with Engenia®, XtendiMax® and Enlist® One.
- Glyphosate alternative for control of actively growing grasses prior to planting.
- In-crop postemergence application.


Application Information:

- Crop oil concentrate (1% v/v) is the preferred adjuvant. Use 2% v/v under arid conditions.
- **Assure II** use rates for non-Enlist volunteer corn control when applied alone:
 - 4 fl oz/A for up to 12-inch corn.
 - 5 fl oz/A for 12- to 18-inch corn.
 - 8 fl oz/A for 18- to 30-inch corn.
- Controls many annual grass weeds using 8 fl oz/A, even in tank mixtures. SEE LABEL FOR SPECIES.


For More Information:

- Contact your local AMVAC® retailer or AMVAC representative to learn more about controlling volunteer corn with **Assure II**, or visit us at AMVAC.com/products/assure-ii.

Volunteer Corn



Volunteer Corn Control, 28 DAA



AMVAC 2020 third party trials 20C01H051-IA402, IN509, ND310, NE311, NE401, and OH 510. Applied to corn 10-12" in height.

Assure II tank mixtures with Enlist One effectively control volunteer corn up to 12" in height at 6 Midwest field trials

See our entire line of products at AMVAC.com

Weeds Controlled and Rate Selection

	Size at Application (in)	Assure II Applied Alone (fl oz product/A)	Assure II Tank-Mixed With Broadleaf Herbicide (fl oz product/A)*
Annual Grasses**			
Volunteer Corn (<i>Zea mays</i>)***	6–30	4–8	8–12
Foxtail, Giant (<i>Setaria faberi</i>)	2–4 (pre-tiller)	5–8	8–12
Johnsongrass, Seedling (<i>Sorghum halepense</i>)	2–8		
Shattercane (<i>Sorghum bicolor</i>)	6–12		
Wild Proso Millet (<i>Panicum miliaceum</i>)	2–6		
Crowfootgrass (<i>Dactyloctenium aegyptium</i>)	2–6	7–8	8–12
Fall Panicum (<i>Panicum dichotomiflorum</i>)	2–6		
Field Sandbur (<i>Cenchrus incertus</i>)	2–6		
Foxtail, Bristly (<i>Setaria verticillata</i>)	2–4		
Foxtail, Giant (<i>Setaria faberi</i>)	2–8		
Foxtail, Green (<i>Setaria viridis</i>)	2–4		
Foxtail, Yellow (<i>Setaria pumila</i>)	2–4		
Goosegrass (<i>Eleusine indica</i>)	2–6†		
Itchgrass (<i>Rottboellia exaltata</i>)	2–8		
Sprangletop (<i>Leptochloa filiformis</i>)	2–6		
Volunteer Barley (<i>Hordeum vulgare</i>)	2–6		
Volunteer Oat (<i>Avena sativa</i>)	2–6		
Volunteer Rye (<i>Secale cereale</i>)	2–6		
Volunteer Wheat (<i>Triticum aestivum</i>)	2–6		
Wild Oat (<i>Avena fatua</i>)	2–6		
Witchgrass (<i>Panicum capillare</i>)	2–6		
Barnyardgrass (<i>Echinochloa crus-galli</i>)	2–6		
Crabgrass, Large (<i>Digitaria sanguinalis</i>)	2–6†		
Crabgrass, Smooth (<i>Digitaria ischaemum</i>)	2–6†		
Junglerice (<i>Echinochloa colonum</i>)	2–6		
Texas Panicum (<i>Panicum texanum</i>)†	2–4		
Red Rice (<i>Oryza sativa</i>)	1–4	9–10	Split†
Woolly Cupgrass (<i>Eriochloa villosa</i>)	2–4§		
Broadleaf Signalgrass (<i>Brachiaria platyphylla</i>)	2–6	10	Split†
Downy Brome (<i>Bromus tectorum</i>)	2–6	10–12	12
Italian Ryegrass (<i>Lolium multiflorum</i>)	2–6		
Jointed Goatgrass (<i>Aegilops cylindrica</i>)	2–6		
Windgrass (<i>Bromus mollis</i>)	2–6		
Perennial Grasses**			
Wirestem Muhly (<i>Muhlenbergia frondosa</i>)	4-8	8–10	Split†
Bermudagrass (<i>Cynodon dactylon</i>)	3 in tall (or up to 6 in runners)	10–12	Split†
Johnsongrass, Rhizome (<i>Sorghum halepense</i>)	10–24		
Quackgrass (<i>Agropyron repens</i>)	6–10		

* Refer to the **Assure II** product label for more information regarding tank mixes with additional herbicides.

** For annual and perennial grasses, up to 12 fl oz/A may be applied based on local recommendations. Under arid conditions, apply at the higher use rate.

*** Does not control quizalofop resistant corn (e.g. Enlist corn). For all other corn, apply 4 fl oz/A of **Assure II** for up to 12-inch tall corn. Apply 5 fl oz/A of **Assure II** for 12- to 18-inch tall corn; use 8 fl oz/A of **Assure II** for 18- to 30-inch tall corn.

† Split Application. Grass species may not be controlled when **Assure II** is tank-mixed with broadleaf herbicides. To prevent antagonism of grass control, apply **Assure II** either 1 day before or 7 days after the broadleaf herbicide. Use rates listed under **Assure II** Applied Alone.

‡ Length of lateral growth.

§ Do not apply **Assure II** at less than 10 fl oz/A in Texas and other areas of the arid west.

§ Size in height or diameter, whichever is more restrictive. Application to plants with more than three tillers may result in unsatisfactory control.