

FLEXIBLE BROADLEAF WEED CONTROL IN SOYBEAN AND FIELD CORN

Product Advantages

- Excellent selectivity fits in both corn and soybean.
- Foundational control of more than 30 broadleaf weeds.*
- Absorption occurs by both roots and shoots.
- Can be tank-mixed with preplant or preemergence herbicides.
- Works in minimum and conventional tillage.

Application Information

(See product label for specific information)

- Preplant incorporated, preplant surface, preemergence in corn and soybean.
 - 1.0–1.33 oz/A (consult product label for recommended rates).
 - Apply up to 30 days before planting.
 - 0.8–1.0 oz/A with triazine-containing pre-mix products for control of velvetleaf and lambsquarters, including triazine-resistant biotypes, in field corn.
- Postemergence
 - Velvetleaf (field corn only) – 0.46–0.93 oz/A from emergence (spike stage) to 20-inch tall corn.
 - Teaweed/Prickly sida control
 - In soybean: 0.125 oz/acre, up to 2-inch sida
 - In field corn: 0.23 – 0.69 oz/acre, up to 6-inch sida
 - Use in AL, AR, DE, GA, KY, LA, MD, MS MO, NJ, NC, OH, PA, SC, TN and VA only

Application Instructions

- Check label for liquid fertilizer mixing or dry fertilizer impregnation instructions.
- Apply in 10–40 gal/A.
- PPI applications incorporate uniformly into top 2–3 inches of soil.
- Check label for pH restrictions and soil insecticide interactions pertaining to corn.
- Rainfast in 6 hours.

**Python will not control ALS-resistant weeds*

Weed Spectrum

(See label for Weed Specific Directions)

Cocklebur ¹	Smooth/Redroot Pigweed
Kochia ²	Sunflower ¹
Lambsquarters	Teaweed (Prickly Sida) ³
Mallow, Venice	Ragweed, Common
Marestail (Horseweed)	Ragweed, Giant
Mustard, Wild	Russian Thistle
Nightshade Spp. ¹	Velvetleaf ³
Sicklepod ¹	Wormwood, Biennial ²
Smartweed, Pennsylvania	

Refer to Python® label for additional weeds controlled

¹Light-moderate Infestations ²Partial Control

³Also Controlled With Postemergence Application

Crop Rotation Interval

Corn, Soybean	0 Months
Alfalfa, Wheat, Barley, Dry Bean and Pea	4 Months
Rice	6 Months
Cotton ¹ , Popcorn and Tobacco	9 Months
Potato, Grain Sorghum	12 Months
Sunflower, Cotton	18 Months
Sugar Beet, Canola	26 Months and Bioassay

¹Following a season maximum Python rate of 0.25 oz/acre

Refer to Python® label for additional crop rotation interval restrictions

See our entire line of products at AMVAC.com

Python Herbicide
HRAC Group 2