



Same great control at half the rate

With AbbA ULTRA® miticide/insecticide from AMVAC, **stone fruit** growers have an innovative tool in the fight against a wide spectrum of damaging mites, including the two-spotted spider mite, Pacific spider mite and European red mite.

AbbA ULTRA is created with the most advanced formulation technology, providing exceptional efficacy while meeting clean air standards. With twice the concentration of abamectin than other formulations, rate **ABBA ULTRA QUICK FACTS:**

Active Ingredient: Abamectin 3.74% Activity: Translaminar movement for extended residual control Mode of Action: Chloride Channel Activators Group Insecticide Number: 6 Signal Word: Warning Formulation: Emulsifiable Concentrate Restricted Use: Yes Packaging Info: 4x1 gal. case; 36 cases/144 gals. = pallet

conversion is easy. Simply use half of the standard .15 abamectin rate. This innovative change with AbbA ULTRA means half the storage, half the transportation and half the container disposal – saving users time and money without sacrificing performance.

An excellent foundational tool in a miticide spray program with rotation partners, AbbA ULTRA provides proven performance at the right price.

PEST	DILUTE SPRAYS ¹ USE RATE	CONCENTRATE SPRAYS ² USE RATE
European red mite	1.25 – 2.5 fl. oz./100 gal.	5 – 10 fl. oz./A
Pacific spider mite	plus	plus
Two-spotted spider mite	0.25% or 1 gal./A horticultural	horticultural spray oil at a
	spray oil or nonionic surfactant at the labeled rate	minimum of 1 gal./A

¹ The rate of AbbA ULTRA per 100 gal. is based on volume of 400 gal./A dilute spray.

To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This can result in use of less than 5 fl. oz./A on small trees.

USE RESTRICTIONS:

Do not apply aerially.

Do not reapply within 21 days of first application.

Do not apply within 21 days of harvest.

Do not apply more than 10 fl. oz./A of AbbA ULTRA or any other foliarly applied abamectin product per application.

Do not apply more than 20 fl. oz./A and/or more than 2 application of AbbA ULTRA or any other foliarly applied abamectin product per cropping season. Do not allow livestock to graze in treated orchards.

For more information about AbbA ULTRA and the entire AMVAC portfolio, please contact your retailer or PCA, visit us online at amvac-chemical.com or call 1-888-GO-AMVAC.

Please consult the full label for crop- and pest-specific recommendations, and follow all label recommendations.