



Same great control at half the rate

With Abba ULTRA[®] miticide/insecticide from AMVAC[®], **cotton** growers have an innovative tool in the fight against a wide spectrum of mites including two-spotted spider mites, Pacific spider mites, strawberry spider mites and carmine spider mites.

Abba ULTRA is created with the most advanced formulation technology, providing exceptional efficacy while meeting clean air standards. With twice the concentration of abamectin than other formulations,

ABBA ULTRA QUICK FACTS:

Active Ingredient: Abamectin 3.74%
 Activity: Translaminar movement for extended residual control
 Mode of Action: Chloride Channel Activators
 Group Insecticide Number: 6
 Signal Word: Warning
 Formulation: Emulsifiable Concentrate
 Restricted Use: Yes
 Packaging Info: 4x1 gal case; 36 cases/144 gal = pallet.

rate conversion is easy. Simply use half of the standard .15 abamectin rate. This innovative change with Abba ULTRA means half the storage, half the transportation and half the container disposal – saving users time and money without sacrificing performance.

An excellent foundational tool in a miticide spray program with rotation partners, Abba ULTRA provides proven performance at the right price.

PEST	EARLY SEASON* USE RATE	IN SEASON USE RATE
Two-spotted spider mite	Apply 2 – 3 fl oz/A when cotton is no more than 10 inches high. Do not use less than 2 fl oz/A	Apply 4 – 8 fl oz/A when mites are first observed and repeat applications as necessary. Use rates depend on the size of the plant and the density of the foliage.
Carmine spider mites		
Pacific spider mites		
Strawberry spider mites		

***West of the Rockies:** the lower use rates can only be used on cotton that is no more than 10 inches high and on applications that are made with ground equipment only.

USE RESTRICTIONS:

Do not apply more than 2 applications of Abba ULTRA or any other foliarly applied abamectin product per cropping season.

Do not apply more than 8 fl oz/A of Abba ULTRA or any other foliarly applied abamectin product per application.

Do not apply more than 16 fl oz/A of Abba ULTRA or any other foliarly applied abamectin product per cropping season.

Do not apply at intervals of less than 21 days.

Do not apply within 20 days of harvest.

Do not feed or allow livestock to graze treated cotton.

For more information about Abba ULTRA and the entire AMVAC cotton portfolio, please contact your retailer, visit us online at AMVAC.com or call 1-888-GO-AMVAC.

Please consult the full label for crop- and pest-specific recommendations, and follow all label recommendations.