

THE RELIABLE SOLUTION FOR FLEAHOPPER CONTROL

When fleahoppers are the problem, Bidrin[®] 8 Insecticide is the solution: Quick knockdown of adult fleahoppers with systemic activity to provide short-term residual control of nymphs. Unlike acephate, Bidrin does not flare spider mites and is the perfect partner in rotation with neonicotinoid insecticides for resistance management.

Bidrin – Key Benefits

- Does not flare spider mites and aphids
- 3-hour rainfast period
- Quick knockdown and systemic control
- Alternative mode of action to neonicotinoid seed treatments
- Tank-mix flexibility
- 3-day REI


Early Season Use

Emergence to pre-bloom protection that stops damage to pinhead and small squares through pre-bloom. A single application of 3.2 fl oz/A of Bidrin is labeled for early season fleahopper control – do not apply more than 3.2 fl oz/A of Bidrin (0.2 lb ai/A of dicotophos) or make more than one application during the early season (emergence to pre-bloom growing stage). The 3.2 fl oz/A rate restriction applies only to early season use.

Bidrin is a valuable tool for mid-to-late season plant bug and stink bug control. Once cotton reaches the first bloom stage of growth, Bidrin may be applied at rates of 4 to 8 fl oz/A. Maximum seasonal use for pest control after first bloom is 16 fl oz/A.



Fleahoppers are piercing-sucking feeders – very similar to tarnished plant bugs. Fleahoppers can cause devastating damage during the first three weeks of squaring and lead to harvest delays and heavy yield loss.



See our entire line of products at AMVAC.com

Bidrin 8 Insecticide
 IRAC Group 1B

