

PLANT BUG CONTROL IN COTTON

Bidrin® 8 Insecticide provides consistent control of piercing-sucking pests. An integral part of plant bug management programs, **Bidrin** delivers the added benefits of superior stink bug control and suppression of aphids.

Key Benefits for Plant Bug Programs

- Quick knockdown of both adults and nymph plant bugs
- Rainfast within 3 hours
- Locally systemic activity for 3 to 5 days residual control
- Excellent tank-mix flexibility
- Suppression of aphids
- Does not flare spider mites
- Superior control of all stink bug species
- Ideal rotation choice in IRM Programs

Late Season Use Information (First Bloom to 30 Days Prior to Harvest)

- Do not apply until “first bloom” stage of cotton
- Suggested use rate 6.0 to 8.0 fl oz/A
- Minimum interval between applications is 14 days
- Do not apply more than 16 fl oz/A of Bidrin (1 lb of dicotophos) during this growth period
- May be applied by ground or air
- REI of 3 days


NOTE: See label for maximum seasonal application rates of dicotophos.



Plant bug adult

Bidrin® 8 Plant Bug Control

Number of Nymphs 6 Days After Treatment
Marianna, AR



See our entire line of products at AMVAC.com

Bidrin 8 Insecticide
IRAC Group 1B

