

GIVE YOUR COTTON A JUMP START BY CONTROLLING THRIPS EARLY FOR TOP YIELD POTENTIAL



Typical crinkling and cupping of leaves caused by thrips damage.

Bidrin[®] 8 Insecticide – Early-Season Control for Strong Stands:

- Does not flare spider mites and aphids
- 3-hour rainfast period
- Quick knockdown and systemic control
- Alternative mode of action to neonicotinoid seed treatments
- Tank-mix flexibility
- 3-day REI

Directions for Early Season Use (Emergence to Pre-Square)


INSECT	DOSAGE (fl oz/A)
Thrips	1.6 to 3.2
Cotton Fleahoppers	1.6 to 3.2
Aphids (suppression)	1.6 to 3.2
Spider Mites (suppression)	1.6 to 3.2
Grasshoppers	3.2

Directions for Application

Do not apply more than 3.2 fl oz/A of Bidrin (0.2 lb a.i./A of dicrotophos) or more than one application during this growth period. Apply by air or ground. Do not graze livestock on treated fields or feed treated gin trash.

Efficacy of Bidrin in Cotton – Thrips

Consultant Demonstration Program Results, 2017 & 2018



See our entire line of products at AMVAC.com

Bidrin 8 Insecticide
IRAC Group 1B

