

# ENHANCE YOUR YELLOW NUTSEDGE AND BROADLEAF WEED CONTROL IN SOYBEAN AND PEANUT

## Product Advantages

- **Classic<sup>®</sup> Herbicide** provides excellent control of many broadleaf weed species\* in soybean and peanut.
- Excellent yellow nutsedge control complements glufosinate and glyphosate herbicides.
- Can be applied on all soybean varieties.
- Excellent tank-mix partner with both preemergence and postemergence herbicides.
- Has both preemergence and postemergence activity.
- Easy to mix formulation.
- Approved tank mix partner with Engenia<sup>®</sup>, Tavium<sup>®</sup>, XtendiMax<sup>®</sup>, Enlist One<sup>®</sup> and Enlist Duo<sup>®</sup>.

## Application Information

### Soybean

- **Preplant (including burndown) and preemergence**  
**Rate:** 1–3 oz/A
- **Postemergence** (2 applications max per season)  
First trifoliolate to 60 days before soybean maturity  
0.33–0.75 oz/A to all soybean varieties
  - Applications up to 1.5 oz/A can be applied to STS/Bolt beans
  - Use 0.5 oz/A minimum for yellow nutsedge control.

### Peanut

- **Postemergence**  
60 days after emergence to 45 days preharvest  
**Rate:** 0.5 oz/A

## Application Instructions

- 15 gal/A
- NIS at 0.25% v/v or COC @ 1% v/v
- UAN at 2 qt/A or AMS @ 2 lb/A
- Rainfast in 1 hour

\*Classic will not control ALS-resistant weed species.

## Control Target Weeds



Common  
Cocklebur<sup>1,2</sup>



Common  
Ragweed<sup>1,2</sup>



Giant  
Ragweed<sup>1</sup>



Sicklepod<sup>1</sup>



Hemp Sesbania<sup>1</sup>



Yellow Nutsedge<sup>1</sup>

Webster, USDA, Bugwood.org  
Schwartz, CSU, Bugwood.org  
Miller and Bodner, SWSS, Bugwood.org

## Weed Spectrum (including those above)

(See label for specific use information)

Burcucumber <sup>3</sup>	Smartweed, ladysthumb <sup>1,2</sup>
Dandelion <sup>1</sup>	Smartweed, Pennsylvania <sup>1,2</sup>
Jimsonweed <sup>1</sup>	Sunflower <sup>1</sup>
Marestail <sup>1,2</sup>	Velvetleaf <sup>1,2</sup>
Morningglory, annual <sup>1</sup>	
Mustard <sup>1</sup>	

<sup>1</sup>Postemergence control

<sup>2</sup>Residual control

<sup>3</sup>Postemergence suppression

## Crop Rotation Interval\*\*

Soybean	0 months
Cereal grains	3 to 4
Field corn	8 to 10
Sweet corn	9 to 18
Dry beans, kidney beans, peas and snap bean	9 to 12
Cotton	8 to 10

\*\*See label for other crops, specific regional rotation intervals,  
use rate and soil parameters.

See our entire line of products at [AMVAC.com](http://AMVAC.com)

**Classic<sup>®</sup> Herbicide**  
HRAC Group 2