POSTEMERGENCE WEED CONTROL IN A GLYPHOSATE-TOLERANT SOYBEAN PROGRAM

Classic® Herbicide Delivers:

- Improved postemergence control of key weeds such as morningglory, hemp sesbania, yellow nutsedge, dandelion and velvetleaf when compared to using glyphosate alone.
- Residual activity to help keep fields clean.
- Excellent tank-mix partner with other postemergence soybean products to expand the weed control spectrum.

Rate Recommendation

• Classic 0.25-0.33 oz/A + glyphosate.

Application Information

- For best control of morningglories and dandelion, a higher rate of Classic is recommended.
- Refer to the glyphosate label for specific ammonium sulfate and adjuvant recommendations.
- Classic may be applied anytime after the first trifoliate has opened but no later than 60 days before soybean maturity.

Weeds Controlled by Classic 0.25–0.33 oz/A¹ + Glyphosate²			
weeds Controlled by Classic 0.25-0.33 02/A* + Glypnosale-			
Weeds	Maximum Height (in inches)	Weeds	Maximum Height (in inches)
Barnyardgrass	6	Panicum: Fall, Texas	10
Cocklebur	8	Pigweed: Redroot, Rough	12
Corn, Volunteer	20	Pigweed, Other	8
Crabgrass Species	10	Ragweed: Common, Giant	8
Dandelion	4	Sesbania, Hemp	4
Foxtail Species	10	Sicklepod	4
Jimsonweed	10	Teaweed (Prickly Sida)	4
Ladysthumb	8	Signalgrass, Broadleaf	4
Lambsquarters	6	Smartweed, Pennsylvania	8
Morningglory: Entireleaf, Ivyleaf, Pitted, Tall	4	Spiderwort, Tropical ³	2
Nightshade, Eastern Black	5	Sunflower	8
Nutsedge, Yellow	6	Velvetleaf	4

¹Classic will not control ALS-resistant weeds.

See our entire line of products at AMVAC.com

Classic Herbicide HRAC Group 2



²Equivalent of 1 qt/A of 4 lb/gal glyphosate.

³Classic at 0.5 oz/A + glyphosate will suppress tropical spiderwort that is no larger than 2 inches in size.