

# ENHANCE YOUR BROADLEAF WEED CONTROL IN SOYBEAN AND PEANUT

## Product Advantages

- Excellent control of many broadleaf weed species in soybean and peanut.
- Can be applied on all soybean varieties.
- Excellent tank-mix partner with both preemergence and postemergence herbicides.
- Has both preemergence and postemergence activity.
- Easy to mix formulation.

## Application Information

### Soybean

- **Preplant (including burndown) and preemergence**  
**Rate:** 1–3 oz/A
- **Postemergence** (2 applications max per season)  
First trifoliolate to 60 days before soybean maturity  
0.33–0.75 oz/A to all soybean varieties  
– Applications up to 1.5 oz/A can be applied to STS/Bolt beans

### Peanut

- **Postemergence**  
60 days after emergence to 45 days preharvest  
**Rate:** 0.5 oz/A

## Application Instructions

- 15 gal/A
- NIS at 0.25% V/V or COC @ 1%V/V
- UAN at 32 oz/A or AMS @ 2 lb/A
- Rainfast in 1 hour

## Control Target Weeds



Common Cocklebur



Common Ragweed



Giant Ragweed



Sicklepod



Hemp Sesbania



Yellow Nutsedge

Webster, USDA, Bugwood.org  
Schwartz, CSU, Bugwood.org  
Miller and Bodner, SWSS, Bugwood.org

## Weed Spectrum

(See label for specific use information)

Burcucumber  
Dandelion  
Morningglory, Annual  
Smartweed, Pennsylvania  
Velvetleaf

## Crop Rotation Interval\*

Soybean	0 months
Cereal grains	3 to 4
Field corn	8 to 10
Sweet corn	9 to 18
Dry beans, kidney beans, peas and snap bean	9 to 12
Cotton	8 to 10

\*See label for other crops, specific regional rotation intervals, use rate and soil parameters.

See our entire line of products at [AMVAC.com](http://AMVAC.com)

**Classic® Herbicide**  
HRAC Group 2