

PREEMERGENCE AND POSTEMERGENCE CONTROL OF KEY BROADLEAF WEEDS IN SOYBEAN

Product Advantages

- Provides effective, flexible and economical preemergence and postemergence control of troublesome broadleaf weeds.
- Can be tank-mixed or applied sequentially with glyphosate to help control glyphosate-tolerant biotypes*.
- Offers freedom of rotation to most major crops and may be applied preplant incorporated, preplant surface, preemergence or postemergence.

Crop Rotation Interval

Wheat	4 Months
Alfalfa, Corn, Cotton, Sorghum, Rice, Dry Beans, Oats, Peanuts	9 Months
Barley	12 Months
Potatoes, Sweet Corn	18 Months
Sugar Beets, Sunflowers	30 Months and Bioassay

Refer to the FirstRate® Herbicide label for additional crop rotation interval restrictions.

Application Timing

- **Preplant/preemergence:** Apply up to two weeks prior to planting or preemergence for additional burndown and residual control.
Rate: 0.3 to 0.6 oz/A
- **Postemergence:** Tank-mix with glyphosate to control tough weeds, including glyphosate-resistant weeds.
FirstRate® Herbicide may be applied any time prior to the R2 (full flower) growth stage of soybeans.
Rate: 0.3 to 0.6 oz/A.
(recommended rate of 0.45 oz)

*FirstRate will not control ALS-resistant weeds.

Control Target Weeds



Lambsquarters



Marestail



Pigweed



Common Ragweed



Giant Ragweed



Morningglory

Weed Spectrum

(See label for specific use information)

Burcucumber ^{2,3}	Marestail (Horseweed)	Sicklepod ²
Cocklebur, Common	Marshelder ²	Smartweed, Pennsylvania
Copperleaf, Hophornbeam ^{2,3}	Morningglory Spp.	Sunflower
Dayflower Spp. ²	Mustard, Wild ²	Thistle, Canada ^{2,3}
Jimsonweed	Pigweed, Redroot/Smooth ¹	Velvetleaf
Lambsquarters ¹	Ragweed, Common	Yellow Nutsedge ^{2,3}
Mallow, Venice	Ragweed, Giant	

Weeds without footnotes 1 and 2 are controlled with both residual and postemergence applications.

¹Residual activity only ²Postemergence only ³Suppression

See our entire line of products at AMVAC.com

FirstRate Herbicide
HRAC Group 2