



# FOLEX<sup>®</sup> 6 EC

## AT THE CORE OF FAST, EFFICIENT COTTON DEFOLIATION PROGRAMS FLEXIBLE • RELIABLE • ECONOMICAL

### Why is Folex<sup>®</sup> the core component of harvest aid programs?

Mature lead drop is a core need in most programs.

Folex<sup>®</sup> 6 EC Cotton Defoliant has excellent tank-mix flexibility to customize defoliation programs based on local conditions and needs.

- Common tank-mix partners are Thidiazuron and ethephon based products.

Folex use rates generally range from 4 to 12 fluid oz/acre depending on tank mix partners and temperature.

- Use higher rates as temperatures decrease.

### Folex<sup>®</sup> 6 EC Cotton Defoliant:

**Adjuvants:** Folex 6 EC may be applied with (1) commercial blends of vegetable or petroleum-based oils, (2) non-ionic surfactant and (3) diesel oil (if allowed by local conditions).

**Rainfastness:** Folex is rainfast one hour after application.

**Plant Response:** Cotton defoliation will start approximately 3 days after application and reach satisfactory levels 5-7 DAT. With low temperatures or drought-stressed conditions, defoliation may extend to 10-14 DAT.

## Folex Based Cotton Harvest Aid Guidelines for the Southeast

Based on University Guidelines from the Southeast

Temperatures	Regrowth Potential	Boll Opening Needed	Number of Passes
<b>HOT</b> Low > 70 degrees; High > 90 degrees	Yes	Yes	One
<b>Folex at 4 to 8 oz/A</b> +thidiazuron at 1.6 to 2.4 oz/A + ethephon at 21 to 24 oz/A or FirstPick™ at 1.5 to 2.0 qts/A			
Temperatures	Regrowth Potential	Boll Opening Needed	Number of Passes
<b>WARM</b> Low > 60degrees; High < 90 Degrees	YES	YES	ONE
<b>Folex at 8 to 12 oz/A</b> +thidiazuron at 1.6 to 2.4 oz/A + ethephon at 24 to 32 oz/A			
Temperatures	Regrowth Potential	Boll Opening Needed	Number of Passes
<b>COOL</b> Low < 60 Degrees; High < 75 Degrees	YES	YES	ONE
<b>Folex at 1 to 1.5 pt/A</b> +thidiazuron* at 1.6 to 2.4 oz/A + ethephon at 1.33 to 2 pt/A or Finish <sup>®</sup> Pro 6 at 26 to 32 oz/A or FirstPick™ at 2 qt.?A *thidiazuron can be dropped if regrowth control is not needed			

If a second application is necessary use rates and products to achieve needed results. Do not exceed seasonal use rates. Folex season use maximum is 24 oz/A.

© 2019 AMVAC Chemical Corporation. All rights reserved. AMVAC, Folex, Yes and Yield Enhancement Solutions are trademarks owned by AMVAC Chemical Corporation. Finish is a registered trademark of Bayer. FirstPick is a trademark of NuFarm. Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or countries. Please check with your local extension service to ensure registration status. All products are EPA registered.

