

POSTEMERGENCE BROADLEAF WEED CONTROL IN FIELD CORN

Product Advantages

- A preferred tank-mix partner with an additional MOA to control difficult weeds such as giant ragweed, dandelion, cocklebur, sunflower and other large seeded broadleaf weeds.
- Use on conventional and all genetically traited field corn.
- Wide application window: Preplant to V6 (6 collar), 20-inch corn.
- Low concern for auxin herbicide volatility.
- Excellent corn tolerance.
- Does not contain atrazine or 2,4-D.

Application Information

Preplant, preemergence and postemergence up to V6 (6 collars) and 20-inch tall corn.

Application Instructions

- 2–5 oz/A (refer to product label for weeds controlled at different rates).
- Use a spray volume of 10–40 gal/A.
- Approved adjuvants required for postemergence application: NIS at 0.25% v/v, COC at 1% v/v, or MSO at 1% v/v plus AMS at 2–4 lb/A or UAN at 2.5% v/v.
- Do not use undiluted UAN as total carrier.

Other Information

- Rainfast in 2 hours.
- Aerial application not allowed.
- Do not apply to sweet corn or popcorn.
- Preharvest interval is 85 days for field corn grown for grain and 45 days for field corn grown for forage or ensilage.
- Maximum total application rate is 6 oz/A in a single crop year.
- Do not apply **Hornet**[®] **Herbicide** if corn was previously treated by the organophosphate insecticides, COUNTER[®] 20G Insecticide or Thimet[®] 20-G Insecticide.

Weed Spectrum (Partial)*

(See label for Weed Specific Directions)

Alfalfa, Volunteer	Mustard Spp.
Burdock	Nightshade Spp.
Canada Thistle	Teaweed (Prickly Sida)
Clover	Purslane
Cocklebur	Ragweed, Common
Curly Dock	Ragweed, Giant
Dandelion	Red Sorrel
Henbit	Sicklepod
Jerusalem Artichoke	Smartweed Spp.
Jimsonweed	Sunflower
Marestail (Horseweed)	Velvetleaf

***Hornet** will not control ALS-resistant weeds that have clopyralid tolerance.

Crop Rotation Interval

Barley, Oat, Rye, Wheat	4 Months
Alfalfa ¹ , Cover Crops ¹ , Dry Beans ¹ , Lima Beans, Forage Grasses ¹ , Popcorn, Rice, Soybean ¹	10.5 Months
Grain Sorghum	12 Months
Snap Bean ¹ , Sweet Corn ¹ , Cotton, Pea ¹ , Peanut, Potato, Sunflower, Tobacco	18 Months
Canola, Sugarbeet and All Other Crops	26 Months and Bioassay

¹Refer to product label for restrictions and different rotation intervals.

See our entire line of products at AMVAC.com

Hornet Herbicide
HRAC Group 2, 4