



PARAZONE®

3SL Herbicide

Parazone® 3SL Herbicide – Now Available from AMVAC®

Key Points about Parazone® 3SL Herbicide from AMVAC:

- Proven 3 lb/gal formulation results in lower product use rates than required for 2 lb/gal formulations
- Excellent mixing characteristics
- Rapidly adsorbed by plant foliage, ParaZone 3SL is rainfast in 30 minutes
- Available in 2 x 2.5 gallon cartons and in 265 gallon mini-bulk totes
- As a Group 22 Herbicide Parazone 3SL has different modes of action than other harvest aid products
- Adjuvant or Crop Oil Concentrate can improve activation of Parazone 3SL. Follow local recommendations.

Application rate ranges allow grower flexibility and fit as a component of most harvest aid programs

- Use at lower rates as a tankmix partner to aid in defoliation and opening of mature bolls during first defoliation pass when planning for sequential applications
- Use at higher specified rates for desiccation of regrowth after an initial defoliation or desiccation application
- Can be used at high rates to aid in controlling regrowth if harvest is delayed

Due to Environmental Variability Follow Local University Recommendations for Use Rates and Tankmix Partners

Use Restrictions — Cotton (Harvest Aid)

- When tank mixing, read and carefully follow all applicable use directions, restrictions, and limitations on the respective product label(s). In interpreting the label(s) of tank mixed products, the most restrictive label limitations must apply.
- Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.
- Because of extremes in environmental and plant conditions, it is advisable to apply the range of rates on a small block of cotton to determine the specific rate that best fits your needs.
- Do not pasture livestock in treated fields or feed treated foliage.
- Do not exceed a total of 32 fl oz (0.75 lb ai) per acre per crop season.
- Up to 4 harvest aid applications may be made per season.
- Allow 7 days between applications.
- Do not apply within 3 days of harvest or grazing.
- Do not harvest within 7 days of application if used with phosphate or chlorate defoliants.
- Do not pasture lactating dairy animals.
- When combined with phosphate defoliants, observe livestock restrictions listed on the phosphate label.

RESTRICTED USE PESTICIDE



©2018 AMVAC Chemical Corporation. AMVAC, and ParaZone 3SL are registered trademarks of AMVAC Chemical Corporation. ParaZone 3SL is a restricted use pesticide. Always read and follow all label directions and use precautions. AV-2018: ParaZone 3SL-Cotton. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.