

FLEXIBLE BROADLEAF WEED CONTROL IN SOYBEANS AND FIELD CORN

Product Features

- Excellent selectivity fits in both corn and soybeans
- Foundational control of more than 30 broadleaf weeds*
- · Absorption occurs by both roots and shoots
- · Can be tank-mixed with preplant or preemergence herbicides
- · Works in minimum and conventional tillage
- Minimal rotation restrictions (refer to product label)

Application Timing

(See product label for specific information)

- Preplant incorporated, preplant surface, preemergence in corn and soubean
 - 1 oz/A (consult product label for recommended rates)
 - · Apply up to 30 days before planting
- Postemergence
 - Velvetleaf (field corn only) 0.46–0.93 oz/A from emergence (spike stage) to 20-inch tall corn
 - Teaweed (prickly sida, soybean only) 0.125 oz/A from first to fifth trifoliate

Application Instructions

- Check label for liquid fertilizer mixing or dry fertilizer impregnation instructions
- Apply in 10–40 gal/A
- PPI applications incorporate uniformly into top 2-3 inches of soil
- Check label for pH restrictions and soil insecticide interactions pertaining to corn

See our entire line of products at **AMVAC.com**

Weed Spectrum

(See label for Weed Specific Directions)

Cocklebur¹ Smooth/Redroot Kochia² Pigweed Lambsquarters Sunflower¹ Mallow, Venice Teaweed (Prickly Sida)³

Marestail
(Horseweed)
Ragweed, Common
Mustard, Wild
Ragweed, Giant
Nightshade Spp.¹
Russian Thistle
Velvetleaf ³

Smartweed, Wormwood, Pennsylvania Biennial ²

Refer to Python® label for additional weeds controlled ¹Light-moderate Infestations ²Partial Control ³Also Controlled With Postemergence Application



Crop Rotation Interval	
Corn, Soybeans	0 Months
Alfalfa, Wheat, Barley, Dry Beans and Peas	4 Months
Rice	6 Months
Potatoes, Grain Sorghum	12 Months
Sunflower, Cotton	18 Months
Sugarbeet, Canola	26 Months and Bioassay

Refer to Python® label for additional crop rotation interval restrictions

Python Herbicide HRAC Group 2



^{*}Python will not control ALS-resistant weeds